

Supplementary Table 3. Correlation between the ratio of proteinuria at 6 months to baseline proteinuria and various cytokines

Cytokine	Control	P value
Serum INF γ (pg/mL)	0.294	0.409
Serum IL-2 (pg/mL)	0.002	0.995
Serum IL-4 (pg/mL)	0.930	< 0.001
Serum IL-10 (pg/mL)	-0.096	0.791
Serum IL-13 (pg/mL)	0.247	0.491
Serum TNF α (pg/mL)	-0.170	0.204
Serum MCP-1 (pg/mL)	-0.269	0.452
Urinary INF γ (pg/mg Cr)	0.189	0.577
Urinary IL-10 (pg/mg Cr)	0.246	0.465
Urinary IL-13 (pg/mg Cr)	0.309	0.356
Urinary MCP-1 (pg/mg Cr)	0.722	0.012
Urinary TNF α (pg/mg Cr)	0.072	0.833
PBMCs INF γ (pg/mL)	0.882	0.002
PBMCs IL-2 (pg/mL)	0.866	0.003
PBMCs IL-4 (pg/mL)	-0.137	0.729
PBMCs IL-10 (pg/mL)	-0.253	0.511
PBMCs IL-13 (pg/mL)	-0.222	0.566
PBMCs MCP-1 (pg/mL)	0.468	0.204
PBMCs TNF α (pg/mL)	0.561	0.112

Data are presented as mean \pm standard deviation.

Cr, creatinine; IgAN, IgA nephropathy; IL, interleukin; INF γ , interferon-gamma; MCP-1, monocyte chemoattractant peptide-1; PBMCs, peripheral blood mononuclear cells; TNF α , tumor necrosis factor alpha.