

Supplementary Table 5. Correlations between the LCx/RCA FAI and continuous covariates in the CKD patients

	LCx FAI		RCA FAI	
	<i>R</i>	p-value	<i>R</i>	p-value
Age	-0.138	0.06	-0.001	0.98
Body mass index	-0.379	<0.001	-0.310	<0.001
Dialysis duration	-0.012	0.89	-0.005	0.95
Hemoglobin	-0.147	0.04	-0.284	<0.001
Urea nitrogen	0.057	0.43	0.144	0.04
Creatinine	0.09	0.21	0.152	0.03
Albumin	-0.199	0.006	-0.235	0.001
Calcium	-0.051	0.49	-0.022	0.75
Phosphorus	0.056	0.44	0.117	0.1
Total cholesterol	-0.176	0.02	-0.140	0.06
Triglyceride	-0.268	0.001	-0.298	<0.001
eGFR	-0.159	0.03	-0.235	0.001
LAD FAI	0.681	<0.001	0.597	<0.001
LCx FAI	1	-	0.692	<0.001
RCA FAI	0.692	<0.001	1	-

The *R* and p-values of correlations between the LCx/RCA FAI and the continuous covariates were calculated using Pearson's correlation coefficient analyses.

eGFR, estimated glomerular filtration rate; FAI, fat attenuation index; LAD, left anterior descending artery; LCx, left circumflex artery; RCA, right coronary artery.