

Supplementary Table 3. Comparison of graft and patient outcomes among kidney transplant recipients with elderly defined as 65 years of age or more

Variable	Old-to-old (n = 16)	Young-to-old (n = 33)	Old-to-young (n = 48)	Young-to-young (n = 926)	p-value
Delayed graft function	1 (6.3)	0 (0)	3 (6.3)	36 (3.9)	0.389
SCr > 1.5 mg/dL at 1 year	4 (25.0)	3 (9.1)	16 (33.3)	133 (14.4)	0.002
Graft failure	0 (0)	1 (3.2)	7 (14.6)	103 (11.1)	0.207
BPAR	0 (0)	2 (6.1)	3 (6.3)	56 (6.0)	0.793
Death	3 (18.8)	7 (21.2)	4 (8.3)	59 (6.4)	0.005
Infection	2 (12.5)	3 (9.1)	2 (4.0)	11 (1.2)	0.001
Cardiovascular disease	0 (0)	1 (3.0)	0 (0)	3 (0.3)	0.19
Cancer	0 (0)	0 (0)	0 (0)	12 (1.3)	>0.999
Others	1 (6.3)	3 (9.1)	2 (4.2)	33 (3.6)	0.211

Data are expressed as number only or number (%).

BPAR, biopsy-proven acute rejection; BPAR, biopsy-proven acute rejection; SCr, serum creatinine.