

Supplementary Table 2. Comparison of predialysis uremic toxins in patients receiving postdilution HDF and MCO dialysis in the first session

Variable	Predialysis in the first session		p-value ^a
	MCO dialysis	Postdilution HDF	
Albumin (mg/dL)	3.7 (3.5–4.0)	3.8 (3.6–3.9)	>0.99
Total protein (mg/dL)	6.6 (6.2–7.1)	6.8 (6.4–7.1)	0.44
BUN (mg/dL)	49.9 (39.8–58.8)	51.4 (42.7–71.8)	0.14
Creatinine (mg/dL)	9.9 (8.6–11.7)	9.5 (9.3–11.1)	0.52
Uric acid (mg/dL)	4.7 (3.3–5.7)	4.9 (3.7–5.6)	0.31
β2MG (μg/L)	23,412 (21,219–25,249)	24,994 (22,778–27,492)	0.07
Myoglobin (ng/mL)	295.0 (211.5–442.0)	267.0 (216.5–399.0)	0.68
Kappa FLC (mg/L)	207.8 (194.0–279.6)	215.9 (202.3–264.3)	0.61
Lambda FLC (mg/L)	163.6 (143.1–218.1)	225.7 (180.8–249.5)	0.06

Data are expressed as medians (interquartile range).

BUN, blood urea nitrogen; β2MG, β2-microglobulin; FLC, free light chain; HDF, hemodiafiltration; MCO, medium cutoff membrane.

^aWilcoxon signed-rank test, p-value for MCO dialysis vs. postdilution HDF.