

Supplementary Table 9. Age-stratified comparative analysis of indications for kidney biopsy

Subgroup	Age (yr)	≤14 yr	15–30 yr	31–45 yr	45–60 yr	>60 yr	Total
AUA	30.87 ± 12.34	100 (6.73)	724 (48.72)	499 (33.58)	121 (8.14)	42 (2.83)	1,486 (14.11)
CNS	38.51 ± 15.95	309 (5.69)	1,607 (29.60)	1,593 (29.34)	1,356 (24.98)	564 (10.39)	5,429 (51.57)
NS	42.48 ± 17.52	104 (3.29)	854 (27.05)	766 (24.26)	880 (27.87)	553 (17.52)	3,157 (29.99)
AKI	41.23 ± 18.86	48 (11.85)	84 (20.74)	86 (21.23)	113 (27.90)	74 (18.27)	405 (3.85)
RPG	47.62 ± 16.54	0 (0)	7 (24.14)	8 (27.59)	11 (37.93)	3 (10.34)	29 (0.28)
Others	30.36 ± 13.67	2 (9.09)	10 (45.45)	5 (22.73)	5 (22.73)	0 (0)	22 (0.21)
Total	38.73 ± 16.53	563 (5.35)	3,286 (31.21)	2,957 (28.09)	2,486 (23.61)	1,236 (11.74)	10,528

Data are expressed as mean ± standard deviation and number (%).

AKI, acute kidney injury; AUA, asymptomatic urinary abnormalities; CNS, chronic nephritic syndrome; NS, nephrotic syndrome; RPGN, rapidly progressive glomerulonephritis.