

Supplementary Table 1. Banff scores and eGFRs in the C4d- negative group

Variable	Absolute ABMR (n = 36)	C4d-positive group (n = 22)	C4d-negative group (n = 4)
MVI score	4.80 ± 1.15	3.91 ± 0.97	3.00 ± 0.80
g	2.24 ± 0.83	1.91 ± 0.75	1.50 ± 0.60
ptc	2.56 ± 0.73	2.00 ± 0.82	1.50 ± 1.00
Other Banff score			
t	1.68 ± 0.94	1.73 ± 1.03	1.83 ± 1.00
i	2.00 ± 0.91	1.73 ± 0.83	2.06 ± 0.50
v	0.28 ± 0.66	0.14 ± 0.48	0.25 ± 0.50
ci	1.11 ± 0.77	1.18 ± 0.85	1.75 ± 0.50
ct	1.00 ± 0.85	1.14 ± 0.88	1.50 ± 1.00
eGFR (mL/min/1.73 m ²)			
3 mo	60.11 ± 14.40	60.55 ± 17.30	78.25 ± 23.90
6 mo	57.19 ± 20.12	54.77 ± 17.40	60.50 ± 18.50
1 yr	60.94 ± 21.78	54.86 ± 21.50	52.00 ± 24.30
2 yr	51.67 ± 19.38	50.05 ± 17.85	46.70 ± 10.40
3 yr	50.52 ± 24.26	44.14 ± 18.96	46.00 ± 11.50
4 yr	41.36 ± 17.85	41.40 ± 18.73	48.00 ± 10.40
5 yr	38.22 ± 21.13	43.18 ± 16.44	45.00 ± 11.40
6 yr	40.82 ± 20.19	41.00 ± 12.23	40.33 ± 11.20

Data are expressed as mean ± standard deviation.

ABMR, antibody-mediated rejection; eGFR, estimated glomerular filtration rate; MVI, microvascular inflammation.