

Supplementary Table 1. Characteristics of pre-KT patients

Clinical index	Non-DM 1	Non-DM 2	Non-DM 3	Non-DM 4	DM 1	DM 2	DM 3	DM 4
Sex	Male	Male	Female	Male	Female	Female	Male	Male
Age (yr)	51	62	42	30	58	57	65	66
HbA1c (%)	5.2	4.2	5.5	6.0	6.2	5.5	5.2	4.5
Hemoglobin (g/dL)	13.7	10.2	7.1	8.5	13.3	8.3	12.9	11.2
Serum Ca (mg/dL)	8.5	8.0	8.8	8.5	9.6	7.7	8.1	9.3
Serum P (mg/dL)	4.3	5.4	5.9	3.6	5.8	3.9	3.7	2.9
Total cholesterol (mg/dL)	163	148	137	145	290	159	67	169
Triglyceride (mg/dL)	71	83	119	55	53	51	46	32
LDL (mg/dL)	102	76	67	94	223	83	28	97
Primary renal disease	IgAN	IgAN	Unknown	IgAN	IgAN	Unknown	Unknown	Unknown
ABO incompatibility	X	X	O	X	X	X	O	X
XM	(-)	(-)	(+)	(-)	(-)	(+)	(-)	(-)
HLA-DSA	(-)	(-)	MFI1808	(-)	(-)	MFI15704	(-)	(-)
Desensitization	(-)	(-)	RTX + PP#6	(-)	(-)	RTX + PP#7	RTX + PP#3	(-)

DM, diabetes mellitus; HbA1c, glycated hemoglobin; HLA-DSA, anti-HLA donor specific antibody; IgAN, IgA nephropathy; KT, kidney transplantation; LDL, low-density lipoprotein; MFI, median fluorescence intensity; PP, plasmapheresis; RTX, rituximab; XM, cross-matching.