

Supplementary Table 2. PSM in PD patient

Characteristic	Before PSM			After PSM		
	COVID-19 (+)	COVID-19 (-)	SMD	COVID-19 (+)	COVID-19 (-)	SMD
No. of patients	98	8,301		98	294	
Age (yr)	56.6 ± 13.7	56.2 ± 13.8	0.026	56.6 ± 13.7	56.1 ± 14.9	0.030
Male sex	58 (59.2)	4,784 (57.6)	0.031	58 (59.2)	174 (59.2)	<0.001
Medical aid	18 (18.4)	1,199 (14.4)	0.106	18 (18.4)	54 (18.4)	<0.001
Metropolitan area	65 (66.3)	4,113 (49.5)	0.345	65 (66.3)	195 (66.3)	<0.001
Vaccination	43 (43.9)	6,319 (76.1)	0.697	43 (43.9)	129 (43.9)	<0.001
Comorbidities						
Diabetes mellitus	30 (30.6)	2,649 (31.9)	0.028	30 (30.6)	81 (27.6)	0.067
Hypertension	41 (41.8)	4,259 (51.3)	0.191	41 (41.8)	117 (39.8)	0.042
Ischemic heart disease	17 (17.3)	878 (10.6)	0.196	17 (17.3)	45 (15.3)	0.055
Heart failure	6 (6.1)	601 (7.2)	0.045	6 (6.1)	21 (7.1)	0.041
Stroke	2 (2.0)	247 (3.0)	0.060	2 (2.0)	8 (2.7)	0.045
Lung disease	3 (3.1)	235 (2.8)	0.014	3 (3.1)	10 (3.4)	0.019
Liver disease	3 (3.1)	251 (3.0)	0.002	3 (3.1)	10 (3.4)	0.019
Malignancy	3 (3.1)	245 (3.0)	0.006	3 (3.1)	9 (3.1)	<0.001
Dementia	1 (1.0)	67 (0.8)	0.022	1 (1.0)	2 (0.7)	0.037

Data are expressed as number only, number (%), or mean ± standard deviation.

COVID-19, coronavirus disease 2019; PD, peritoneal dialysis; PSM, propensity score matching; SMD, standardized mean difference.