

Supplementary Table 10. HRs for 28- and 90-day mortality based on NPS quartiles

Mortality	Model 1 ^a		Model 2 ^b		Model 3 ^c	
	HR (95% CI)	p-value	HR (95% CI)	p-value	HR (95% CI)	p-value
28-Day						
Q1	Reference		Reference		Reference	
Q2	0.70 (0.59–0.84)	<0.001	0.69 (0.58–0.83)	<0.001	0.60 (0.51–0.72)	<0.001
Q3	0.70 (0.59–0.84)	<0.001	0.69 (0.57–0.82)	<0.001	0.53 (0.44–0.63)	<0.001
Q4	0.72 (0.60–0.86)	<0.001	0.70 (0.59–0.84)	<0.001	0.44 (0.37–0.53)	<0.001
90-Day						
Q1	Reference		Reference		Reference	
Q2	0.74 (0.62–0.87)	<0.001	0.73 (0.62–0.87)	<0.001	0.59 (0.50–0.69)	<0.001
Q3	0.74 (0.63–0.88)	<0.001	0.73 (0.62–0.87)	<0.001	0.52 (0.44–0.62)	<0.001
Q4	0.73 (0.62–0.87)	<0.001	0.72 (0.60–0.85)	<0.001	0.45 (0.38–0.54)	<0.001

CI, confidence interval; HR, hazard ratio; PAR, platelet-to-albumin ratio.

^aUnadjusted model; ^badjusted for age and sex; ^cmodel with additional adjustments for Charlson Comorbidity Index score, and body mass index.