

Supplementary Table 3. KDQOL-36 scores and effect size^a according to suicidal ideation

Patient-reported mental health	Suicidal ideation		Effect size Cohen's <i>d</i>
	No (n = 1,563)	Yes (n = 316)	
KD-symptoms and problems	88 ± 12	76 ± 16	1.00 (L)
KD-work	68 ± 38	44 ± 39	0.63 (M)
KD-sexual function	48 ± 46	34 ± 41	0.30 (S)
KD-physician encouragement	82 ± 20	78 ± 20	0.20 (S)
KD-effect of kidney disease	90 ± 12	77 ± 18	1.08 (L)
KD-cognitive function	92 ± 11	80 ± 17	1.09 (L)
KD-sleep	74 ± 16	64 ± 18	0.63 (M)
KD-patient satisfaction	63 ± 22	57 ± 21	0.27 (S)
KD-burden of kidney disease	71 ± 25	47 ± 29	0.96 (L)
KD-quality of social interaction	78 ± 15	64 ± 20	0.93 (L)
KD-social support	74 ± 22	55 ± 24	0.86 (L)
SF-physical function	88 ± 15	77 ± 21	0.73 (M)
SF-role physical	85 ± 30	64 ± 39	0.70 (M)
SF-pain	86 ± 19	72 ± 25	0.74 (M)
SF-general health	44 ± 21	29 ± 19	0.71 (M)
SF-emotional well-being	67 ± 16	47 ± 18	1.25 (L)
SF-role emotional	86 ± 30	59 ± 44	0.90 (L)
SF-social function	89 ± 14	71 ± 24	1.29 (L)
SF-energy/fatigue	53 ± 18	35 ± 18	1.00 (L)
Component summary			
KDCS	75 ± 11	62 ± 13	1.18 (L)
PCS (physical)	76 ± 16	60 ± 20	1.00 (L)
MCS (mental)	73 ± 15	53 ± 21	1.33 (L)

Data are expressed as mean ± standard deviation.

KD, kidney disease; KDCS, kidney disease component summary; KD-QOL-36, Kidney Disease Quality of Life 36-Item; MCS, mental component summary; PCS, physical component summary; SF, short form.

^aEffect size (standardized mean difference) is considered; 0.2 to 0.5 small (S), 0.5 to 0.8 moderate (M), and >0.8 large (L).