

Supplementary Table 1. Comparisons of body mass index and body surface area according to type of SDKT and immunological risk

Complication	Standard-risk group			High-risk group		
	W2H (n = 291)	H2W (n = 75)	p-value	W2H (n = 34)	H2W (n = 55)	p-value
Body mass index (kg/m ²)						
Donor	23.3 ± 3.1	24.8 ± 2.7	<0.001**	23.4 ± 2.8	24.6 ± 2.4	0.03*
Recipient	24.2 ± 3.1	22.1 ± 3.7	<0.001**	23.5 ± 2.5	22.2 ± 3.8	0.07
R/D	1.1 ± 0.2	0.9 ± 0.2	<0.001**	1.2 ± 0.1	0.9 ± 0.1	0.01*
Body surface area (m ²)						
Donor	1.6 ± 0.1	1.8 ± 0.1	<0.001**	1.6 ± 0.1	1.8 ± 0.1	<0.001**
Recipient	1.8 ± 0.1	1.5 ± 0.1	<0.001**	1.8 ± 0.1	1.6 ± 0.1	<0.001**
R/D	1.1 ± 0.1	0.9 ± 0.1	<0.001**	1.0 ± 0.1	0.9 ± 0.2	<0.001**

Data are expressed as mean ± standard deviation.

H2W, husband-to-wife recipient; R/D, recipient-to-donor ratio; SDKT, spousal donor kidney transplantation; W2H, wife-to-husband recipient.

*p < 0.05 and **p < 0.01 by Student t-test.