

**Supplementary Table 2. Baseline characteristics of C3G patients based on ESKD progression**

Characteristic	Progression to ESKD (n = 8)	Non progression to ESKD (n = 34)	p-value
Age (yr)	64.0 ± 20.5	42.5 ± 21.0	0.01
Male sex			
Weight (kg)	60.8 ± 11.7	62.2 ± 13.2	0.81
SBP (mmHg)	145.9 ± 29.3	131.0 ± 23.8	0.14
DBP (mmHg)	75.5 ± 12.1	79.4 ± 17.1	0.55
eGFR (mL/min/1.73 m <sup>2</sup> )	24.3 ± 17.17.2	62.6 ± 39.6	0.01
Creatinine (mg/dL)	5.3 ± 6.3	2.0 ± 2.2	0.01
BUN (mg/dL)	44.7 ± 30.0	27.6 ± 20.5	0.08
Hemoglobin (g/L)	10.3 ± 1.5	11.9 ± 2.3	0.17
Protein (g/dL)	5.7 ± 0.9	6.3 ± 0.9	0.15
Albumin (g/dL)	3.1 ± 0.4	3.3 ± 0.8	0.58
Cholesterol (mg/dL)	199.8 ± 87.7	182.5 ± 60.7	0.52
CRP (mg/dL)	1.2 ± 1.5	1.3 ± 2.5	0.93
UPCR (g/g)	3.6 ± 3.0	2.5 ± 2.6	0.33
UACR (g/g) <sup>a</sup>	3.1 ± 2.8	0.7 ± 0.5	0.02
Hypertension	6 (75.0)	19 (57.6)	0.45
Diabetes mellitus	2 (28.6)	2 (6.7)	0.16
Coronary artery disease	1 (14.3)	1 (3.4)	0.36
Cardiovascular disease	1 (14.3)	4 (13.8)	>0.99

Data are expressed as mean ± standard deviation or number (%).

BUN, blood urea nitrogen; CRP, C-reactive protein; C3G, complement 3 glomerulopathy; DBP, diastolic blood pressure; eGFR, estimated glomerular filtration rate; ESKD, end-stage kidney disease; SBP, systolic blood pressure; UACR, urine albumin-to-creatinine ratio; UPCR, urine protein-to-creatinine ratio.

<sup>a</sup>UACR was measured in 169 patients.