

Supplementary Table 1. STOP-BANG score distribution according to questionnaire items in by group

Variable	Non-albuminuria					Albuminuria				
	Total	0-2	3-4	5-8	p-value	Total	0	3-4	5-8	p-value
No. of patients	9,857	5,119	3,666	1,072		1,066	358	495	213	
Snoring (%)	20.12 ± 0.53	3.77 ± 0.31	25.88 ± 0.91	71.61 ± 1.59	<0.001	24.19 ± 1.67	4.67 ± 1.48	17.01 ± 1.91	69.34 ± 3.74	<0.001
Tired (%)	29.48 ± 0.59	19.60 ± 0.64	33.48 ± 0.94	58.90 ± 1.79	<0.001	31.89 ± 1.59	11.66 ± 1.97	35.46 ± 2.49	54.68 ± 4.02	<0.001
Observed (%)	9.85 ± 0.36	0.78 ± 0.14	8.71 ± 0.55	51.99 ± 1.71	<0.001	11.39 ± 1.30	0.62 ± 0.44	6.20 ± 1.33	39.00 ± 4.29	<0.001
HTN, pressure (%)	35.24 ± 0.65	17.97 ± 0.63	46.89 ± 1.00	72.09 ± 1.63	<0.001	67.52 ± 1.80	46.49 ± 3.05	73.58 ± 2.48	86.10 ± 3.05	<0.001
BMI (%)	5.38 ± 0.31	1.06 ± 0.18	6.41 ± 0.52	20.51 ± 1.49	<0.001	8.42 ± 1.11	2.52 ± 1.21	6.81 ± 1.24	20.85 ± 3.50	<0.001
Age (%)	69.09 ± 0.79	59.65 ± 0.95	77.81 ± 1.07	81.79 ± 1.46	<0.001	83.53 ± 1.50	75.57 ± 2.95	86.63 ± 2.00	88.77 ± 2.91	0.001
Neck (%)	40.98 ± 0.55	9.26 ± 0.50	66.78 ± 0.95	94.75 ± 0.75	<0.001	49.92 ± 1.86	6.56 ± 1.81	60.91 ± 2.53	91.52 ± 2.49	<0.001
Gender, male (%)	48.71 ± 0.45	18.55 ± 0.65	75.43 ± 0.79	92.97 ± 0.88	<0.001	53.05 ± 1.86	9.22 ± 2.06	66.53 ± 2.41	89.95 ± 2.45	<0.001

Data are expressed as number only or mean ± standard error.

BMI, body mass index; HTN, hypertension.